

Fixed points and fuzzy stability of an additive-quadratic functional equation

Choonkil Park¹, Abbas Najati² and Sun Young Jang^{3*}

¹Department of Mathematics, Research Institute for Natural Sciences,
Hanyang University, Seoul 133-791, Korea,
e-mail: baak@hanyang.ac.kr

²Department of Mathematical Sciences,
University of Mohaghegh Ardabili, Ardabil 56199-11367, Iran
e-mail: a.nejati@yahoo.com

³Department of Mathematics, University of Ulsan, Ulsan 680-749, Korea
e-mail: jsym@uou.ulsan.ac.kr

Abstract. The fuzzy stability problems for the Cauchy additive functional equation and the Jensen additive functional equation in fuzzy Banach spaces have been investigated by Moslehian et al.

Using fixed point method, we prove the Hyers-Ulam stability of the functional equation

$$\begin{aligned} lf\left(\sum_{i=1}^l x_i\right) + \sum_{i=1}^l f\left(lx_i - \sum_{j=1}^l x_j\right) & \quad (0.1) \\ & = \frac{l^2+l}{2} \sum_{i=1}^l f(x_i) + \frac{l^2-l}{2} \sum_{i=1}^l f(-x_i) \quad (l \geq 2) \end{aligned}$$

in fuzzy Banach spaces.

Keywords: fuzzy Banach space, fixed point, functional equation associated with inner product space, Hyers-Ulam stability, quadratic mapping, additive mapping.

1. INTRODUCTION AND PRELIMINARIES

Katsaras [?] defined a fuzzy norm on a vector space to construct a fuzzy vector topological structure on the space. Some mathematicians have defined fuzzy norms on a vector space from various points of view [?, ?, ?]. In particular, Bag and Samanta [?], following Cheng and Mordeson [?], gave an idea of fuzzy norm in such a manner that the corresponding fuzzy metric is of Kramosil and Michalek type [?]. They established a decomposition theorem of a fuzzy norm into a family of crisp norms and investigated some properties of fuzzy normed spaces [?].

We use the definition of fuzzy normed spaces given in [?, ?, ?] to investigate a fuzzy version of the Hyers-Ulam stability for the functional equation (0.1) in the fuzzy normed vector space setting.

Definition 1.1. [?, ?, ?, ?] Let X be a real vector space. A function $N : X \times \mathbb{R} \rightarrow [0, 1]$ is called a *fuzzy norm* on X if for all $x, y \in X$ and all $s, t \in \mathbb{R}$,

- (N₁) $N(x, t) = 0$ for $t \leq 0$;
- (N₂) $x = 0$ if and only if $N(x, t) = 1$ for all $t > 0$;
- (N₃) $N(cx, t) = N(x, \frac{t}{|c|})$ if $c \neq 0$;
- (N₄) $N(x + y, s + t) \geq \min\{N(x, s), N(y, t)\}$;

⁰2010 Mathematics Subject Classification: Primary 46S40; 46C05; 39B52; 46S50; 26E50; 47H10.

*Corresponding author.

(N₅) $N(x, \cdot)$ is a non-decreasing function of \mathbb{R} and $\lim_{t \rightarrow \infty} N(x, t) = 1$;

(N₆) for $x \neq 0$, $N(x, \cdot)$ is continuous on \mathbb{R} .

The pair (X, N) is called a *fuzzy normed vector space*.

The properties of fuzzy normed vector spaces and examples of fuzzy norms are given in [?, ?].

Definition 1.2. [?, ?, ?, ?] Let (X, N) be a fuzzy normed vector space. A sequence $\{x_n\}$ in X is said to be *convergent* or *converge* if there exists an $x \in X$ such that $\lim_{n \rightarrow \infty} N(x_n - x, t) = 1$ for all $t > 0$. In this case, x is called the *limit* of the sequence $\{x_n\}$ and we denote it by $N\text{-}\lim_{n \rightarrow \infty} x_n = x$.

Definition 1.3. [?, ?, ?] Let (X, N) be a fuzzy normed vector space. A sequence $\{x_n\}$ in X is called *Cauchy* if for each $\varepsilon > 0$ and each $t > 0$ there exists an $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ and all $p > 0$, we have $N(x_{n+p} - x_n, t) > 1 - \varepsilon$.

It is well-known that every convergent sequence in a fuzzy normed vector space is Cauchy. If each Cauchy sequence is convergent, then the fuzzy norm is said to be *complete* and the fuzzy normed vector space is called a *fuzzy Banach space*.

We say that a mapping $f : X \rightarrow Y$ between fuzzy normed vector spaces X and Y is continuous at a point $x_0 \in X$ if for each sequence $\{x_n\}$ converging to x_0 in X , then the sequence $\{f(x_n)\}$ converges to $f(x_0)$. If $f : X \rightarrow Y$ is continuous at each $x \in X$, then $f : X \rightarrow Y$ is said to be *continuous* on X (see [?]).

The stability problem of functional equations originated from a question of Ulam [?] concerning the stability of group homomorphisms. Hyers [?] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [?] for additive mappings and by Th.M. Rassias [?] for linear mappings by considering an unbounded Cauchy difference. A generalization of the Th.M. Rassias theorem was obtained by Găvruta [?] by replacing the unbounded Cauchy difference by a general control function in the spirit of Th.M. Rassias' approach.

The functional equation

$$f(x + y) + f(x - y) = 2f(x) + 2f(y)$$

is called a *quadratic functional equation*. In particular, every solution of the quadratic functional equation is said to be a *quadratic function*. A Hyers-Ulam stability problem for the quadratic functional equation was proved by Skof [?] for mappings $f : X \rightarrow Y$, where X is a normed space and Y is a Banach space. Cholewa [?] noticed that the theorem of Skof is still true if the relevant domain X is replaced by an Abelian group. Czerwik [?] proved the Hyers-Ulam stability of the quadratic functional equation. The stability problems of several functional equations have been extensively investigated by a number of authors and there are many interesting results concerning this problem (see [?]-[?], [?, ?], [?]-[?], [?]-[?]).

In [?], the author proved that if an even mapping $f : V \rightarrow W$ satisfies the functional equation (0.1), then the even mapping $f : V \rightarrow W$ is quadratic, and that if an odd mapping $f : V \rightarrow W$ satisfies the functional equation (0.1), then the odd mapping $f : V \rightarrow W$ is Cauchy additive. Moreover, the author proved the Hyers-Ulam stability of the quadratic functional equation (0.1) in real Banach spaces.

Let X be a set. A function $d : X \times X \rightarrow [0, \infty]$ is called a *generalized metric* on X if d satisfies

- (1) $d(x, y) = 0$ if and only if $x = y$;
- (2) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (3) $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in X$.

We recall a fundamental result in fixed point theory.

Theorem 1.4. [?, ?] *Let (X, d) be a complete generalized metric space and let $J : X \rightarrow X$ be a strictly contractive mapping with Lipschitz constant $L < 1$. Then for each given element $x \in X$, either*

$$d(J^n x, J^{n+1} x) = \infty$$

for all nonnegative integers n or there exists a positive integer n_0 such that

- (1) $d(J^n x, J^{n+1} x) < \infty, \quad \forall n \geq n_0;$
- (2) *the sequence $\{J^n x\}$ converges to a fixed point y^* of J ;*
- (3) y^* *is the unique fixed point of J in the set $Y = \{y \in X \mid d(J^{n_0} x, y) < \infty\}$;*
- (4) $d(y, y^*) \leq \frac{1}{1-L} d(y, Jy)$ *for all $y \in Y$.*

In 1996, G. Isac and Th.M. Rassias [?] were the first to provide applications of stability theory of functional equations for the proof of new fixed point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [?], [?], [?], [?], [?], [?]).

This paper is organized as follows: In Section 2, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for an odd case. In Section 3, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for an even case.

Throughout this paper, assume that X is a vector space and that (Y, N) is a fuzzy Banach space. Let l be a fixed integer greater than 1.

2. HYERS-ULAM STABILITY OF THE FUNCTIONAL EQUATION (0.1): AN ODD CASE

In this section, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for an odd case.

For a given mapping $f : X \rightarrow Y$, we define

$$\begin{aligned} Cf(x_1, \dots, x_l) : &= lf \left(\sum_{i=1}^l x_i \right) + \sum_{i=1}^l f \left(lx_i - \sum_{j=1}^l x_j \right) \\ &- \frac{l^2 + l}{2} \sum_{i=1}^l f(x_i) - \frac{l^2 - l}{2} \sum_{i=1}^l f(-x_i) \end{aligned}$$

for all $x_1, \dots, x_l \in X$.

Using fixed point method, we prove the Hyers-Ulam stability of the functional equation $Cf(x_1, \dots, x_l) = 0$ in fuzzy Banach spaces: an odd case.

Theorem 2.1. *Let $\varphi : X^l \rightarrow [0, \infty)$ and $\psi(x) := \underbrace{\varphi(x, \dots, x)}_{l \text{ times}}$ be functions such that there exists an $L < 1$ with $\varphi(x_1, \dots, x_l) \leq \frac{L}{l} \varphi(lx_1, \dots, lx_l)$ for all $x_1, \dots, x_l \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying*

$$N(Cf(x_1, \dots, x_l), t) \geq \frac{t}{t + \varphi(x_1, \dots, x_l)} \tag{2.1}$$

for all $x_1, \dots, x_l \in X$ and all $t > 0$. Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} l^n f\left(\frac{x}{l^n}\right)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(l^2 - l^2 L)t}{(l^2 - l^2 L)t + L\psi(x)} \tag{2.2}$$

for all $x \in X$ and all $t > 0$.

Proof. Letting $x_1 = \cdots = x_l = x$ in (2.1), we get

$$N(lf(lx) - l^2f(x), t) \geq \frac{t}{t + \underbrace{\varphi(x, \dots, x)}_{l \text{ times}}} = \frac{t}{t + \psi(x)} \quad (2.3)$$

for all $x \in X$.

Consider the set

$$S := \{g : X \rightarrow Y\}$$

and introduce the generalized metric on S :

$$d(g, h) = \inf\{\mu \in \mathbb{R}_+ : N(g(x) - h(x), \mu t) \geq \frac{t}{t + \psi(x)}, \forall x \in X, \forall t > 0\},$$

where, as usual, $\inf \phi = +\infty$. It is easy to show that (S, d) is complete. (See the proof of Lemma 2.1 of [?].)

Now we consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := lg\left(\frac{x}{l}\right)$$

for all $x \in X$.

Let $g, h \in S$ be given such that $d(g, h) = \varepsilon$. Then

$$N(g(x) - h(x), \varepsilon t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. Hence

$$\begin{aligned} N(Jg(x) - Jh(x), L\varepsilon t) &= N\left(lg\left(\frac{x}{l}\right) - lh\left(\frac{x}{l}\right), L\varepsilon t\right) \\ &= N\left(g\left(\frac{x}{l}\right) - h\left(\frac{x}{l}\right), \frac{L}{l}\varepsilon t\right) \\ &\geq \frac{\frac{L}{l}\varepsilon t}{\frac{L}{l}\varepsilon t + \psi\left(\frac{x}{l}\right)} \geq \frac{\frac{L}{l}\varepsilon t}{\frac{L}{l}\varepsilon t + \frac{L}{l}\psi(x)} \\ &= \frac{t}{t + \psi(x)} \end{aligned}$$

for all $x \in X$ and all $t > 0$. So $d(g, h) = \varepsilon$ implies that $d(Jg, Jh) \leq L\varepsilon$. This means that

$$d(Jg, Jh) \leq Ld(g, h)$$

for all $g, h \in S$.

It follows from (2.3) that

$$N\left(f(x) - lf\left(\frac{x}{l}\right), \frac{Lt}{l^2}\right) \geq \frac{\frac{L}{l}t}{\frac{L}{l}t + \psi\left(\frac{x}{l}\right)} \geq \frac{t}{t + \psi(x)} \quad (2.4)$$

for all $x \in X$. So $d(f, Jf) \leq \frac{L}{l^2}$.

By Theorem 1.4, there exists a mapping $A : X \rightarrow Y$ satisfying the following:

(1) A is a fixed point of J , i.e.,

$$A\left(\frac{x}{l}\right) = \frac{1}{l}A(x) \quad (2.5)$$

for all $x \in X$. Since $f : X \rightarrow Y$ is odd, $A : X \rightarrow Y$ is an odd mapping. The mapping A is a unique fixed point of J in the set

$$M = \{g \in S : d(f, g) < \infty\}.$$

Fuzzy stability of additive-quadratic functional equation

This implies that A is a unique mapping satisfying (2.5) such that there exists a $\mu \in (0, \infty)$ satisfying

$$N(f(x) - A(x), \mu t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$;

(2) $d(J^n f, A) \rightarrow 0$ as $n \rightarrow \infty$. This implies the equality

$$N\text{-}\lim_{n \rightarrow \infty} l^n f\left(\frac{x}{l^n}\right) = A(x)$$

for all $x \in X$;

(3) $d(f, A) \leq \frac{1}{1-L}d(f, Jf)$, which implies the inequality

$$d(f, A) \leq \frac{L}{l^2 - l^2 L}.$$

This implies that the inequality (2.2) holds.

By (2.1),

$$N\left(l^n C f\left(\frac{x_1}{l^n}, \dots, \frac{x_l}{l^n}\right), l^n t\right) \geq \frac{t}{t + \varphi\left(\frac{x_1}{l^n}, \dots, \frac{x_l}{l^n}\right)}$$

for all $x_1, \dots, x_l \in X$, all $t > 0$ and all $n \in \mathbb{N}$. So

$$N\left(l^n C f\left(\frac{x_1}{l^n}, \dots, \frac{x_l}{l^n}\right), t\right) \geq \frac{\frac{t}{l^n}}{\frac{t}{l^n} + \frac{L^n}{l^n} \varphi(x_1, \dots, x_l)}$$

for all $x_1, \dots, x_l \in X$, all $t > 0$ and all $n \in \mathbb{N}$. Since $\lim_{n \rightarrow \infty} \frac{\frac{t}{l^n}}{\frac{t}{l^n} + \frac{L^n}{l^n} \varphi(x_1, \dots, x_l)} = 1$ for all $x_1, \dots, x_l \in X$ and all $t > 0$,

$$N(CA(x_1, \dots, x_l), t) = 1$$

for all $x_1, \dots, x_l \in X$ and all $t > 0$. Thus the mapping $A : X \rightarrow Y$ is additive, as desired. \square

Corollary 2.2. Let $\theta \geq 0$ and let p be a real number with $p > 1$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an odd mapping satisfying

$$N(Cf(x_1, \dots, x_l), t) \geq \frac{t}{t + \theta \sum_{j=1}^l \|x_j\|^p} \quad (2.6)$$

for all $x_1, \dots, x_l \in X$ and all $t > 0$. Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} l^n f\left(\frac{x}{l^n}\right)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(l^p - l)t}{(l^p - l)t + \theta \|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 2.1 by taking

$$\varphi(x_1, \dots, x_l) := \theta \sum_{j=1}^l \|x_j\|^p$$

for all $x_1, \dots, x_l \in X$. Then we can choose $L = l^{1-p}$ and we get the desired result. \square

Theorem 2.3. Let $\varphi : X^l \rightarrow [0, \infty)$ and $\psi(x) := \underbrace{\varphi(x, \dots, x)}_{l \text{ times}}$ be functions such that there exists an $L < 1$ with $\varphi(x_1, \dots, x_l) \leq lL\varphi\left(\frac{x_1}{l}, \dots, \frac{x_l}{l}\right)$ for all $x_1, \dots, x_l \in X$. Let $f : X \rightarrow Y$ be an odd mapping

satisfying (2.1). Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{l^n} f(l^n x)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(l^2 - l^2 L)t}{(l^2 - l^2 L)t + \psi(x)} \quad (2.7)$$

for all $x \in X$ and all $t > 0$.

Proof. Let (S, d) be the generalized metric space defined in the proof of Theorem 2.1.

Consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := \frac{1}{l}g(lx)$$

for all $x \in X$.

Let $g, h \in S$ be given such that $d(g, h) = \varepsilon$. Then

$$N(g(x) - h(x), \varepsilon t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. Hence

$$\begin{aligned} N(Jg(x) - Jh(x), L\varepsilon t) &= N\left(\frac{1}{l}g(lx) - \frac{1}{l}h(lx), L\varepsilon t\right) \\ &= N(g(lx) - h(lx), lL\varepsilon t) \\ &\geq \frac{lLt}{lLt + \psi(lx)} \geq \frac{lLt}{lLt + lL\psi(x)} \\ &= \frac{t}{t + \psi(x)} \end{aligned}$$

for all $x \in X$ and all $t > 0$. So $d(g, h) = \varepsilon$ implies that $d(Jg, Jh) \leq L\varepsilon$. This means that

$$d(Jg, Jh) \leq Ld(g, h)$$

for all $g, h \in S$.

It follows from (2.3) that

$$N\left(f(x) - \frac{1}{l}f(lx), \frac{t}{l^2}\right) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. Thus $d(f, Jf) \leq \frac{1}{l^2}$.

By Theorem 1.4, there exists a mapping $A : X \rightarrow Y$ satisfying the following:

(1) A is a fixed point of J , i.e.,

$$A(lx) = lA(x) \quad (2.8)$$

for all $x \in X$. Since $f : X \rightarrow Y$ is odd, $A : X \rightarrow Y$ is an odd mapping. The mapping A is a unique fixed point of J in the set

$$M = \{g \in S : d(f, g) < \infty\}.$$

This implies that A is a unique mapping satisfying (2.8) such that there exists a $\mu \in (0, \infty)$ satisfying

$$N(f(x) - A(x), \mu t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$;

(2) $d(J^n f, A) \rightarrow 0$ as $n \rightarrow \infty$. This implies the equality

$$N\text{-}\lim_{n \rightarrow \infty} \frac{1}{l^n} f(l^n x) = A(x)$$

for all $x \in X$;

(3) $d(f, A) \leq \frac{1}{1-L}d(f, Jf)$, which implies the inequality

$$d(f, A) \leq \frac{1}{l^2 - l^2L}.$$

This implies that the inequality (2.7) holds.

The rest of the proof is similar to the proof of Theorem 2.1. \square

Corollary 2.4. *Let $\theta \geq 0$ and let p be a real number with $0 < p < 1$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.6). Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{l^n} f(l^n x)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that*

$$N(f(x) - A(x), t) \geq \frac{(l - l^p)t}{(l - l^p)t + \theta \|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 2.3 by taking

$$\varphi(x_1, \dots, x_l) := \theta \sum_{j=1}^l \|x_j\|^p$$

for all $x_1, \dots, x_l \in X$. Then we can choose $L = l^{p-1}$ and we get the desired result. \square

3. HYERS-ULAM STABILITY OF THE FUNCTIONAL EQUATION (0.1): AN EVEN CASE

In this section, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for an even case.

Using fixed point method, we prove the Hyers-Ulam stability of the functional equation $Cf(x_1, \dots, x_l) = 0$ in fuzzy Banach spaces: an even case.

Theorem 3.1. *Let $\varphi : X^l \rightarrow [0, \infty)$ and $\psi(x) := \underbrace{\varphi(x, \dots, x)}_{l \text{ times}}$ be functions such that there exists an $L < 1$ with $\varphi(x_1, \dots, x_l) \leq \frac{L}{l^2} \varphi(lx_1, \dots, lx_l)$ for all $x_1, \dots, x_l \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.1). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} l^{2n} f\left(\frac{x}{l^n}\right)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that*

$$N(f(x) - Q(x), t) \geq \frac{(l^3 - l^3L)t}{(l^3 - l^3L)t + L\psi(x)} \quad (3.1)$$

for all $x \in X$ and all $t > 0$.

Proof. Letting $x_1 = \dots = x_l = x$ in (2.1), we get

$$N(lf(lx) - l^3f(x), t) \geq \frac{t}{t + \underbrace{\varphi(x, \dots, x)}_{l \text{ times}}} = \frac{t}{t + \psi(x)} \quad (3.2)$$

for all $x \in X$.

Let (S, d) be the generalized metric space defined in the proof of Theorem 2.1.

Now we consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := l^2g\left(\frac{x}{l}\right)$$

for all $x \in X$.

Let $g, h \in S$ be given such that $d(g, h) = \varepsilon$. Then

$$N(g(x) - h(x), \varepsilon t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. Hence

$$\begin{aligned} N(Jg(x) - Jh(x), L\epsilon t) &= N\left(l^2g\left(\frac{x}{l}\right) - l^2h\left(\frac{x}{l}\right), L\epsilon t\right) \\ &= N\left(g\left(\frac{x}{l}\right) - h\left(\frac{x}{l}\right), \frac{L}{l^2}\epsilon t\right) \geq \frac{\frac{Lt}{l^2}}{\frac{Lt}{l^2} + \psi\left(\frac{x}{l}\right)} \\ &\geq \frac{\frac{Lt}{l^2}}{\frac{Lt}{l^2} + \frac{L}{l^2}\psi(x)} = \frac{t}{t + \psi(x)} \end{aligned}$$

for all $x \in X$ and all $t > 0$. So $d(g, h) = \epsilon$ implies that $d(Jg, Jh) \leq L\epsilon$. This means that

$$d(Jg, Jh) \leq Ld(g, h)$$

for all $g, h \in S$.

It follows from (3.2) that

$$N\left(f(x) - l^2f\left(\frac{x}{l}\right), \frac{Lt}{l^3}\right) \geq \frac{\frac{Lt}{l^2}}{\frac{Lt}{l^2} + \psi\left(\frac{x}{l}\right)} \geq \frac{t}{t + \psi(x)} \quad (3.3)$$

for all $x \in X$. So $d(f, Jf) \leq \frac{L}{l^3}$.

By Theorem 1.4, there exists a mapping $Q : X \rightarrow Y$ satisfying the following:

(1) Q is a fixed point of J , i.e.,

$$Q\left(\frac{x}{l}\right) = \frac{1}{l^2}Q(x) \quad (3.4)$$

for all $x \in X$. Since $f : X \rightarrow Y$ is even, $Q : X \rightarrow Y$ is an even mapping. The mapping Q is a unique fixed point of J in the set

$$M = \{g \in S : d(f, g) < \infty\}.$$

This implies that Q is a unique mapping satisfying (3.4) such that there exists a $\mu \in (0, \infty)$ satisfying

$$N(f(x) - Q(x), \mu t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$;

(2) $d(J^n f, Q) \rightarrow 0$ as $n \rightarrow \infty$. This implies the equality

$$N\text{-}\lim_{n \rightarrow \infty} l^{2n} f\left(\frac{x}{l^n}\right) = Q(x)$$

for all $x \in X$;

(3) $d(f, Q) \leq \frac{1}{1-L}d(f, Jf)$, which implies the inequality

$$d(f, Q) \leq \frac{L}{l^3 - l^3 L}.$$

This implies that the inequality (3.1) holds.

The rest of the proof is similar to the proof of Theorem 2.1. □

Corollary 3.2. *Let $\theta \geq 0$ and let p be a real number with $p > 2$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.6). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} l^{2n} f\left(\frac{x}{l^n}\right)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that*

$$N(f(x) - Q(x), t) \geq \frac{(l^p - l^2)t}{(l^p - l^2)t + \theta\|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 3.1 by taking

$$\varphi(x_1, \dots, x_l) := \theta \sum_{j=1}^l \|x_j\|^p$$

for all $x_1, \dots, x_l \in X$. Then we can choose $L = l^{2-p}$ and we get the desired result. \square

Similarly, we can obtain the following. We will omit the proof.

Theorem 3.3. Let $\varphi : X^l \rightarrow [0, \infty)$ and $\psi(x) := \varphi(\underbrace{x, \dots, x}_{l \text{ times}})$ be functions such that there exists an $L < 1$ with $\varphi(x_1, \dots, x_l) \leq l^2 L \varphi(\frac{x_1}{l}, \dots, \frac{x_l}{l})$ for all $x_1, \dots, x_l \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.1). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{l^{2n}} f(l^n x)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(l^3 - l^3 L)t}{(l^3 - l^3 L)t + \psi(x)}$$

for all $x \in X$ and all $t > 0$.

Corollary 3.4. Let $\theta \geq 0$ and let p be a real number with $0 < p < 2$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.6). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{l^{2n}} f(l^n x)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that

$$N(f(x) - Q(x), t) \geq \frac{(l^2 - l^p)t}{(l^2 - l^p)t + \theta \|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 3.3 by taking

$$\varphi(x_1, \dots, x_l) := \theta \sum_{j=1}^l \|x_j\|^p$$

for all $x_1, \dots, x_l \in X$. Then we can choose $L = l^{p-2}$ and we get the desired result. \square

ACKNOWLEDGMENTS

C. Park was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (NRF-2012R1A1A2004299), and S. Y. Jang was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (NRF-2011-004872) and the Ulsan Metropolitan Office of Education.

REFERENCES

- [1] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2** (1950), 64–66.
- [2] T. Bag and S.K. Samanta, *Finite dimensional fuzzy normed linear spaces*, J. Fuzzy Math. **11** (2003), 687–705.
- [3] T. Bag and S.K. Samanta, *Fuzzy bounded linear operators*, Fuzzy Sets and Systems **151** (2005), 513–547.
- [4] L. Cădariu and V. Radu, *Fixed points and the stability of Jensen's functional equation*, J. Inequal. Pure Appl. Math. **4**, no. 1, Art. ID 4 (2003).

- [5] L. Cădariu and V. Radu, *On the stability of the Cauchy functional equation: a fixed point approach*, Grazer Math. Ber. **346** (2004), 43–52.
- [6] L. Cădariu and V. Radu, *Fixed point methods for the generalized stability of functional equations in a single variable*, Fixed Point Theory and Applications **2008**, Art. ID 749392 (2008).
- [7] S.C. Cheng and J.M. Mordeson, *Fuzzy linear operators and fuzzy normed linear spaces*, Bull. Calcutta Math. Soc. **86** (1994), 429–436.
- [8] P.W. Cholewa, *Remarks on the stability of functional equations*, Aequationes Math. **27** (1984), 76–86.
- [9] S. Czerwik, *On the stability of the quadratic mapping in normed spaces*, Abh. Math. Sem. Univ. Hamburg **62** (1992), 59–64.
- [10] P. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific Publishing Company, New Jersey, Hong Kong, Singapore and London, 2002.
- [11] J. Diaz and B. Margolis, *A fixed point theorem of the alternative for contractions on a generalized complete metric space*, Bull. Amer. Math. Soc. **74** (1968), 305–309.
- [12] M. Eshaghi Gordji, *A characterization of (σ, τ) -derivations on von Neumann algebras*, Politehn. Univ. Bucharest Sci. Bull. Ser. A–Appl. Math. Phys. **73** (2011), No. 1, 111–116.
- [13] M. Eshaghi Gordji, A. Bodaghi and C. Park, *A fixed point approach to the stability of double Jordan centralizers and Jordan multipliers on Banach algebras*, Politehn. Univ. Bucharest Sci. Bull. Ser. A–Appl. Math. Phys. **73** (2011), No. 2, 65–74.
- [14] M. Eshaghi Gordji, H. Khodaei and R. Khodabakhsh, *General quartic-cubic-quadratic functional equation in non-Archimedean normed spaces*, Politehn. Univ. Bucharest Sci. Bull. Ser. A–Appl. Math. Phys. **72** (2010), No. 3, 69–84.
- [15] M. Eshaghi Gordji and M.B. Savadkouhi, *Approximation of generalized homomorphisms in quasi-Banach algebras*, An. Stiint. Univ. Ovidius Constanta Ser. Mat. **17** (2009), No. 2, 203–213.
- [16] C. Felbin, *Finite dimensional fuzzy normed linear spaces*, Fuzzy Sets and Systems **48** (1992), 239–248.
- [17] Z. Gajda, *On stability of additive mappings*, Internat. J. Math. Math. Sci. **14** (1991), 431–434.
- [18] P. Găvruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **184** (1994), 431–436.
- [19] D.H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [20] D.H. Hyers, G. Isac and Th.M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [21] G. Isac and Th.M. Rassias, *Stability of ψ -additive mappings: Applications to nonlinear analysis*, Internat. J. Math. Math. Sci. **19** (1996), 219–228.
- [22] S. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press Inc., Palm Harbor, Florida, 2001.
- [23] A.K. Katsaras, *Fuzzy topological vector spaces II*, Fuzzy Sets and Systems **12** (1984), 143–154.
- [24] I. Kramosil and J. Michalek, *Fuzzy metric and statistical metric spaces*, Kybernetika **11** (1975), 326–334.
- [25] S.V. Krishna and K.K.M. Sarma, *Separation of fuzzy normed linear spaces*, Fuzzy Sets and Systems **63** (1994), 207–217.
- [26] D. Mihet and V. Radu, *On the stability of the additive Cauchy functional equation in random normed spaces*, J. Math. Anal. Appl. **343** (2008), 567–572.
- [27] M. Mirzavaziri and M.S. Moslehian, *A fixed point approach to stability of a quadratic equation*, Bull. Braz. Math. Soc. **37** (2006), 361–376.
- [28] A.K. Mirmostafae, M. Mirzavaziri and M.S. Moslehian, *Fuzzy stability of the Jensen functional equation*, Fuzzy Sets and Systems **159** (2008), 730–738.
- [29] A.K. Mirmostafae and M.S. Moslehian, *Fuzzy versions of Hyers-Ulam-Rassias theorem*, Fuzzy Sets and Systems **159** (2008), 720–729.

- [30] A.K. Mirmostafae and M.S. Moslehian, *Fuzzy approximately cubic mappings*, Inform. Sci. **178** (2008), 3791–3798.
- [31] C. Park, *Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras*, Fixed Point Theory and Applications **2007**, Art. ID 50175 (2007).
- [32] C. Park, *Generalized Hyers-Ulam-Rassias stability of quadratic functional equations: a fixed point approach*, Fixed Point Theory and Applications **2008**, Art. ID 493751 (2008).
- [33] C. Park, *Inner product spaces and quadratic functional equations*, (preprint).
- [34] C. Park, Y. Cho and M. Han, *Functional inequalities associated with Jordan-von Neumann type additive functional equations*, J. Inequal. Appl. **2007**, Art. ID 41820 (2007).
- [35] C. Park and J. Cui, *Generalized stability of C^* -ternary quadratic mappings*, Abstract Appl. Anal. **2007**, Art. ID 23282 (2007).
- [36] C. Park and A. Najati, *Homomorphisms and derivations in C^* -algebras*, Abstract Appl. Anal. **2007**, Art. ID 80630 (2007).
- [37] V. Radu, *The fixed point alternative and the stability of functional equations*, Fixed Point Theory **4** (2003), 91–96.
- [38] Th.M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [39] Th.M. Rassias, *Problem 16; 2*, Report of the 27th International Symp. on Functional Equations, Aequationes Math. **39** (1990), 292–293; 309.
- [40] Th.M. Rassias, *On the stability of the quadratic functional equation and its applications*, Studia Univ. Babeş-Bolyai **XLIII** (1998), 89–124.
- [41] Th.M. Rassias, *The problem of S.M. Ulam for approximately multiplicative mappings*, J. Math. Anal. Appl. **246** (2000), 352–378.
- [42] Th.M. Rassias, *On the stability of functional equations in Banach spaces*, J. Math. Anal. Appl. **251** (2000), 264–284.
- [43] Th.M. Rassias, *On the stability of functional equations and a problem of Ulam*, Acta Appl. Math. **62** (2000), 23–130.
- [44] Th.M. Rassias and P. Šemrl, *On the behaviour of mappings which do not satisfy Hyers-Ulam stability*, Proc. Amer. Math. Soc. **114** (1992), 989–993.
- [45] Th.M. Rassias and P. Šemrl, *On the Hyers-Ulam stability of linear mappings*, J. Math. Anal. Appl. **173** (1993), 325–338.
- [46] Th.M. Rassias and K. Shibata, *Variational problem of some quadratic functionals in complex analysis*, J. Math. Anal. Appl. **228** (1998), 234–253.
- [47] F. Skof, *Proprietà locali e approssimazione di operatori*, Rend. Sem. Mat. Fis. Milano **53** (1983), 113–129.
- [48] S. M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publ. New York, 1960.
- [49] J.Z. Xiao and X.H. Zhu, *Fuzzy normed spaces of operators and its completeness*, Fuzzy Sets and Systems **133** (2003), 389–399.